

Liitetaulukko 1. Vastasyntyneen hypoglykemian hoitorajoja					
		Plasman glukoosipitoisuus, mmol/l			
Suosituksen antaja	Ikä 0–4 tuntia	Ikä 4–24 tuntia	Ikä 24–48 tuntia	Ikä 48–72 tuntia	Ikä > 72 tuntia
American Academy of Pediatrics (Yhdysvallat) ¹	1,4–2,2	2,0–2,5	2,5		
Pediatric Endocrine Society (Eurooppa, Yhdysvallat) ²	2,8			3,3	
Canadian Paediatric Society (Kanada) ³	2,6				2,8
Kansallinen suositus, asiantuntijatyöryhmä (Ruotsi) ⁴	2,6–3,0				
¹ Committee on Fetus and Newborn, Adamkin DH. Postnatal glucose homeostasis in late-preterm and term infants. <i>Pediatr</i> 2011;127:575–9. ² Thornton PS, Stanley CA, De Leon DD, ym. Recommendations from the pediatric endocrine society for evaluation and management of persistent hypoglycemia in neonates, infants, and children. <i>J Pediatr</i> 2015;167:238–45. ³ Narvey MR, Marks SD. The screening and management of newborns at risk for low blood glucose. <i>Paediatr Child Health</i> 2019;24:536–54. ⁴ Wackernagel D, Gustafsson A, Edstedt Bonamy AK ym. Swedish national guideline for prevention and treatment of neonatal hypoglycaemia in newborn infants with gestational age ≥35 weeks. <i>Acta Paediatr</i> 2020;109:31–44.					